

For patients with multiple sclerosis (MS)

LEARNING ABOUT **TREATMENT OPTIONS FOR
MS EXACERBATIONS**



H.P. **Acthar**[®] GEL
(repository corticotropin injection) 80 U/mL





Introduction

There is currently no cure for MS and, for many, living with this condition can be challenging. MS affects you and the people who care about you and it can change the way you live. But, it is manageable and, with proper treatment, most people will lead full and productive lives.

Your doctor may have prescribed a disease-modifying drug (DMD). DMDs are designed to slow the progression of MS, but you still may experience exacerbations of your symptoms.

This brochure includes general information about MS exacerbations and explores treatment options available for you—specifically, you will learn more about Acthar. Take a moment to read this information to help you start a conversation with your doctor or nurse about your treatment.

What is an MS exacerbation?

An exacerbation (also known as a relapse) is a sudden worsening of 1 or more MS symptoms or the appearance of new symptoms. To be considered a new exacerbation, it must last at least 24 hours and be separated from a previous exacerbation by at least 1 month. Sometimes, exacerbations can last anywhere from days to weeks, but may also last for months.

A true MS exacerbation is caused by an area of inflammation in the central nervous system. The inflammation damages the myelin (white, fatty substance that covers and protects a part of nerve cells), and slows or disrupts the transmission of nerve signals, which causes the symptoms of MS.

What are the signs and symptoms of an MS exacerbation?

Symptoms can vary from person to person and some are more common than others. Below are some of the symptoms that may indicate an MS exacerbation; however, this is not a complete list. If you are unsure if a symptom you are experiencing is a sign of an exacerbation or if you have a symptom that is worrying you, talk to your doctor.

- Numbness, weakness, feeling constantly tired
- Loss of muscle function (especially in your lower torso)
- Difficulty going to the bathroom
- Uncontrollable shaking (tremors or muscle spasms)
- Problems with walking or balance
- Difficulty speaking
- Problems with memory and attention
- Changes in your ability to see

Over the course of MS, some symptoms may come and go. Talk to your doctor or nurse if you experience any of the symptoms and about possible treatment options.

What treatments are available for MS exacerbations?

Occasionally, your doctor may need to prescribe medications other than your DMD to help manage a sudden relapse or exacerbation. Acthar and corticosteroids are effective treatments used to successfully treat and control MS exacerbations.

Most patients receive corticosteroids through intravenous (given through a vein) infusion treatments on an outpatient basis, either at a clinic or hospital or at home. In some cases, hospitalization may be recommended or required. Corticosteroids are typically administered by multiple intravenous infusions over a 3- to 5-day course of treatment. High-dose oral steroids are also sometimes used.

While corticosteroids are considered the standard for treating MS exacerbations, corticosteroids may not be right for everyone. Acthar is an effective, flexible treatment option for patients who have unique treatment needs.

What is Acthar?

H.P. Acthar Gel, or Acthar, is one of the first effective medications approved by the FDA for the treatment of acute exacerbations in patients with MS. Acthar is a prescription medication containing the hormone adrenocorticotropin (“a-DRE-no-cor-ti-co-tro-pin”), also known as ACTH.

Acthar has been proven to speed recovery from MS exacerbations. It works by helping the body produce natural steroid hormones to reduce inflammation.

Unlike intravenous corticosteroids, Acthar can be used when and where it is most convenient for you. It can be self-injected or given by a friend, family member, care partner, or your doctor or nurse. Your doctor will advise you on how to inject Acthar and what the appropriate dose is for you.

Who is Acthar for?

Acthar is another treatment option when corticosteroids are not right for you.

Acthar may be the right choice if you:

- Are unable to cope with the side effects of corticosteroids
- Have been treated with a corticosteroid and it did not work
- Don't have the time or access to receive intravenous therapy
- Have trouble receiving medication through your veins

All drugs have the possibility of causing some side effects. Whether you are taking a DMD, corticosteroid, or Acthar, talk to your doctor about the possible side effects associated with your medication.

Please see page 6 for Important Safety Information.



Important Safety Information

Patients, parents, and caregivers should be aware of the important information about Acthar.

Acthar should never be given intravenously (into a vein). Acthar should not be used in patients with a skin condition called scleroderma, bone density loss (osteoporosis), infection throughout the body, eye infection called ocular herpes simplex, recent surgery, history of or a current stomach ulcer, heart problems, high blood pressure, or allergy to pig-derived proteins. Tell your doctor about any health problems or medicines.

Acthar may cause side effects similar to side effects that happen due to treatment with steroid medicines. Not all of these side effects have occurred with Acthar but they may occur. Acthar is a medicine that affects a patient's immune system, and therefore patients may be more likely to get new infections, or inactive infections may become active. Acthar has effects on the adrenal gland. When a patient is taking Acthar, their adrenal gland may produce too much of a hormone called cortisol. This can cause symptoms of Cushing's syndrome (upper body fat, rounded face, thin skin), which is more common in patients who take this medicine for a long time. When a patient stops taking Acthar after a long time, the body may not produce enough cortisol on its own (adrenal insufficiency). The doctor may prescribe a steroid medicine to protect the body until the adrenal gland recovers. Do not stop administering Acthar without talking to your doctor first. Blood pressure should be checked during treatment, and your doctor may instruct you to make some dietary changes. Patients should not receive certain vaccines during treatment with Acthar. Talk to your doctor about which vaccines are safe for use. Acthar may hide (or mask) symptoms of other conditions or diseases and it may be more difficult for your doctor to diagnose other conditions or diseases in you or your child during treatment. The person receiving Acthar has an increased risk for bleeding from the stomach or having a stomach ulcer. Inform your doctor about any pain in the stomach area, bloody vomit, or bloody or black stools. While on Acthar changes in mood and behavior such as irritability, depression, or trouble sleeping may occur.

Other side effects are possible. Acthar may make certain other medical conditions worse, such as diabetes (may increase blood sugar); cause eye problems, such as cataracts, increased pressure in the eye (glaucoma), and possible damage to the optic nerve; and cause allergic reactions to Acthar (seen as skin rash, swelling of the face, tongue, lips, or throat, and trouble breathing). Acthar may affect growth and physical development after long term use. Long term use of Acthar may cause an increase in the size of the heart, but this condition usually goes away after Acthar is stopped.

The most common side effects of Acthar in infants include: infections, increased blood pressure, irritability and changes in behavior, changes in appetite and weight, diarrhea, and vomiting. Other adverse reactions reported in adults and children over 2 years of age included: abdominal bloating, anxiety, asthma, chest discomfort, congestive heart failure, dizziness, shortness of breath, redness of the face, fluid retention, flushing, headache, injection site pain, tiredness, muscle weakness, nervousness, rapid heart rate, and lack of energy. Tell your doctor if there is any side effect that bothers you or your child or that does not go away. These are not all of the possible side effects of Acthar. For more information, ask your doctor, nurse or pharmacist, go to www.acthar.com, or call **1-800-465-9217**. You may report side effects to the FDA at 1-800-FDA-1088.

Please refer to the full Prescribing Information for a description of Acthar.

Patient support for Acthar

Questcor, the maker of Acthar, is committed to helping MS patients get started with Acthar. We know that getting access and paying for MS treatments can be a burden at times, especially if you are on multiple treatments. That is why Questcor started the Acthar Support & Access Program (A.S.A.P.). Created to honor this commitment, the Acthar Support & Access Program is staffed with reimbursement experts who help you in several ways:

- Questcor makes sure cost is not a problem
 - 90% of patients have had Acthar paid for by their insurance company*
 - Over \$50 million has been given in free Acthar†
 - Patient costs are typically less than \$50 for Acthar
 - 100% copay assistance for all patients in need
- Questcor will also help make sure you can get Acthar quickly
 - Time from prescription to treatment is approximately 2.5–4 days‡ (time may vary)
- Questcor provides home injection training support
 - Free easy-to-use, multimedia home injection training materials
 - Coordinates free Home Injection Training Services by licensed nurses

Your doctor will need to complete a prescription form and fax it to the Acthar Support & Access Program. To learn more about the Acthar Support & Access Program, visit www.acthar.com or call **1-888-435-2284**.

Learn more about Acthar

If you are interested in learning more about Acthar, you can:

- Ask your doctor or nurse
- Visit www.acthar.com

Your doctor can describe the benefits and possible side effects associated with Acthar in more detail. He or she can also answer questions about the dosing and convenient self-administration of Acthar. Your doctor will help you determine if Acthar is the right choice for you.

*Based on Acthar referral data from January to August 2010. Does not include those that were withdrawn prior to completion by physician/patient.

†\$50 million in free product has been distributed to patients across all Acthar indications.

‡Measured in cases where drug was shipped and doctor-patient response to A.S.A.P. were timely.

References:

Data on file: RD-001-00. Questcor Pharmaceuticals, Inc.

Exacerbations. National Multiple Sclerosis Society Web site. <http://www.nationalmssociety.org/about-multiple-sclerosis/treatments/exacerbations/index.aspx>. Accessed November 9, 2010.

H.P. Acthar® Gel (repository corticotropin injection) prescribing information, Questcor Pharmaceuticals, Inc.

National Multiple Sclerosis Society. The disease-modifying drugs: newly diagnosed. <http://nationalmssociety.org/multimedia-library/brochures/newly-diagnosed/download.aspx?id=45>. Published October 2008. Accessed November 9, 2010.

Symptoms. National Multiple Sclerosis Society Web site. <http://www.nationalmssociety.org/about-multiple-sclerosis/symptoms/index.aspx>. Accessed November 9, 2010.



**ACTHAR SUPPORT
& ACCESS PROGRAM**
A.S.A.P.

H.P. **Acthar**® GEL
(repository corticotropin injection) 80 U/mL

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use H.P. Acthar Gel safely and effectively. See full prescribing information for H.P. Acthar Gel.

H.P. Acthar Gel (repository corticotropin injection) INJECTION, GEL for INTRAMUSCULAR | SUBCUTANEOUS use

Initial U.S. Approval: 1952

RECENT MAJOR CHANGES

- Indications and Usage, (1) 10/10
- Dosage and Administration, (2) 10/10
- Contraindications, Infantile Spasms (4) 10/10
- Warnings and Precautions (5) 10/10

INDICATIONS AND USAGE

- H.P. Acthar Gel is an adrenocorticotropic hormone (ACTH) analogue indicated as monotherapy for the treatment of infantile spasms in infants and children under 2 years of age. (1.1)
- H.P. Acthar Gel is indicated for the treatment of exacerbations of multiple sclerosis in adults. (1.2)
- H.P. Acthar Gel may be used for the following disorders and diseases: rheumatic; collagen; dermatologic; allergic states; ophthalmic; respiratory; and edematous state; (1.3 to 1.9)

DOSAGE AND ADMINISTRATION

- In the treatment of infantile spasms, the recommended dose is 150 U/m² divided into twice daily intramuscular injections of 75 U/m². After 2 weeks of treatment, dosing should be gradually tapered and discontinued over a 2-week period. (2.1)
- In the treatment of acute exacerbations of multiple sclerosis, daily intramuscular or subcutaneous doses of 80-120 units for 2-3 weeks may be administered. It may be necessary to taper the dose. (2.2)
- In the treatment of other disorders and diseases, dosing will need to be individualized depending on the disease under treatment and the medical condition of the patient. It may be necessary to taper the dose. (2.3)

DOSAGE FORMS AND STRENGTHS

- 5 mL multi-dose vial containing 80 USP units per mL (3)

CONTRAINDICATIONS

- H.P. Acthar Gel should never be given intravenously.
- H.P. Acthar Gel is contraindicated in patients with scleroderma, osteoporosis, systemic fungal infections, ocular herpes simplex, recent surgery, history of or the presence of a peptic ulcer, congestive heart failure, uncontrolled hypertension, or sensitivity to proteins of porcine origin.
- Administration of live or live attenuated vaccines is contraindicated in patients receiving immunosuppressive doses of H.P. Acthar Gel.
- H.P. Acthar Gel is contraindicated in children under 2 years of age with suspected congenital infections. (4)
- Treatment of conditions listed within the INDICATIONS section is contraindicated when they are accompanied by primary adrenocortical insufficiency or adrenocortical hyperfunction. (4)

WARNINGS AND PRECAUTIONS

- Infections: Increased susceptibility to new infection and increased risk of exacerbation, dissemination or reactivation of latent infections. Signs and symptoms of infection may be masked. (5.1)
- Adrenal Insufficiency after Prolonged Therapy: Monitor for effects of hypothalamic-pituitary-axis suppression after stopping treatment. (5.2)

- Cushing's Syndrome: May occur after prolonged therapy. Monitor for signs and symptoms. (5.2)
- Elevated Blood Pressure, Salt and Water Retention and Hypokalemia: Monitor blood pressure and sodium and potassium levels. (5.3)
- Vaccination: Do not administer live or attenuated vaccines to patients on immunosuppressive doses. (5.4)
- Masking of Symptoms of Other Underlying Disease/Disorders. Monitor patients for signs of other underlying disease/disorders that may be masked. (5.5)
- Gastrointestinal Perforation and Bleeding: There is a risk for gastric ulcers and bleeding. There is an increased risk of perforation in patients with certain GI disorders. Signs and symptoms may be masked. Monitor for signs of perforation and bleeding. (5.6)
- Behavioral and Mood Disturbances: May include euphoria, insomnia, mood swings, personality changes, severe depression and psychosis. Existing conditions may be aggravated (5.7)
- Comorbid Diseases: Symptoms of diabetes and myasthenia gravis may be worsened with treatment. (5.8)
- Ophthalmic Effects: Monitor for cataracts, infections and glaucoma. (5.9)
- Immunogenicity Potential: Neutralizing antibodies with chronic administration may lead to a loss of endogenous ACTH activity. (5.10)
- Use in Patients with Hypothyroidism or Liver Cirrhosis: May result in an enhanced effect. (5.11)
- Negative Effects on Growth and Physical Development: Monitor pediatric patients on long term therapy. (5.12)
- Decrease in Bone Density: Monitor for osteoporosis in patients on long term therapy. (5.13)
- Use in Pregnancy: Embryocidal effect. Apprise women of potential harm to the fetus. (5.14)

ADVERSE REACTIONS

- Common adverse reactions for H.P. Acthar Gel are similar to those of corticosteroids and include fluid retention, alteration in glucose tolerance, elevation in blood pressure, behavioral and mood changes, increased appetite and weight gain. (6)
- Specific adverse reactions resulting from drug use in children under 2 years of age are increased risk of infections, hypertension, irritability, Cushingoid symptoms, cardiac hypertrophy and weight gain. (6.1.1)

To report SUSPECTED ADVERSE REACTIONS, contact Questcor Pharmaceuticals, Inc. at (800) 411-3065 or (510) 400-0700 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

DRUG INTERACTIONS

- H.P. Acthar Gel may accentuate the electrolyte loss associated with diuretic therapy. (7)

USE IN SPECIFIC POPULATIONS

- Pregnancy: H.P. Acthar Gel has been shown to have an embryocidal effect and should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. (8.1)
- Pediatric Use: Prolonged use of H.P. Acthar Gel in children may inhibit skeletal growth. If use is necessary, it should be given intermittently with careful observation. (5.12 and 8.3)

See 17 for Patient Counseling Information and FDA-approved Medication Guide

FULL PRESCRIBING INFORMATION: CONTENTS*

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

- 1.1 Infantile spasms:
- 1.2 Multiple Sclerosis:
- 1.3 Rheumatic Disorders:
- 1.4 Collagen Diseases:
- 1.5 Dermatologic Diseases:
- 1.6 Allergic States:
- 1.7 Ophthalmic Diseases:
- 1.8 Respiratory Diseases:
- 1.9 Edematous State:

2 DOSAGE AND ADMINISTRATION

- 2.1 Specific Recommended Dosage Regimen for Infantile Spasms in Infants and Children Under 2 Years of Age
- 2.2 Recommended Dosage Regimen for the Treatment of Acute Exacerbations in Adults with Multiple Sclerosis.
- 2.3 Recommended Dosage Regimen for Other Indications for Adults and Children Over 2 Years of Age
- 2.4 Preparation
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
 - 5.1 Infections

5.2	Cushing's Syndrome and Adrenal Insufficiency Upon Withdrawal
5.3	Elevated Blood Pressure, Salt and Water Retention and Hypokalemia
5.4	Vaccination
5.5	Masking Symptoms of Other Diseases
5.6	Gastrointestinal Perforation and Bleeding
5.7	Behavioral and Mood Disturbances
5.8	Comorbid Diseases
5.9	Ophthalmic Effects
5.10	Immunogenicity Potential
5.11	Use in Patients with Hypothyroidism or Liver Cirrhosis
5.12	Negative Effects on Growth and Physical Development
5.13	Decrease in Bone Density
5.14	Use in Pregnancy
6	ADVERSE REACTIONS
6.1	Clinical Studies Experience
6.1.1	Adverse Reactions in Infants and Children Under 2 Years of Age
6.2	Postmarketing Experience
6.2.1	Allergic Reactions
6.2.2	Cardiovascular
6.2.3	Dermatologic
6.2.4	Endocrine
6.2.5	Gastrointestinal

6.2.6	Metabolic
6.2.7	Musculoskeletal
6.2.8	Neurological
6.3	Possible Additional Steroidogenic Effects
6.3.1	Dermatologic
6.3.2	Endocrine
6.3.3	Metabolic
6.3.4	Musculoskeletal
6.3.5	Neurological
6.3.6	Ophthalmic
7	DRUG INTERACTIONS
8	USE IN SPECIFIC POPULATIONS
8.1	Pregnancy
8.3	Nursing Mothers
8.4	Pediatric Use
10	OVERDOSAGE
11	DESCRIPTION
12	CLINICAL PHARMACOLOGY
12.1	Mechanism of Action
13	NONCLINICAL TOXICOLOGY
13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
14	CLINICAL STUDIES
16	HOW SUPPLIED / STORAGE AND HANDLING
17	PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Infantile spasms:

H.P. Acthar Gel (repository corticotropin injection) is indicated as monotherapy for the treatment of infantile spasms in infants and children under 2 years of age.

1.2 Multiple Sclerosis:

H.P. Acthar Gel (repository corticotropin injection) is indicated for the treatment of acute exacerbations of multiple sclerosis in adults. Controlled clinical trials have shown H.P. Acthar Gel to be effective in speeding the resolution of acute exacerbations of multiple sclerosis. However, there is no evidence that it affects the ultimate outcome or natural history of the disease.

1.3 Rheumatic Disorders:

As adjunctive therapy for short-term administration (to tide the patient over an acute episode or exacerbation) in: Psoriatic arthritis, Rheumatoid arthritis, including juvenile rheumatoid arthritis (selected cases may require low-dose maintenance therapy), Ankylosing spondylitis.

1.4 Collagen Diseases:

During an exacerbation or as maintenance therapy in selected cases of: systemic lupus erythematosus, systemic dermatomyositis (polymyositis).

1.5 Dermatologic Diseases:

Severe erythema multiforme, Stevens-Johnson syndrome.

1.6 Allergic States:

Serum sickness.

1.7 Ophthalmic Diseases:

Severe acute and chronic allergic and inflammatory processes involving the eye and its adnexa such as: keratitis, iritis, iridocyclitis, diffuse posterior uveitis and choroiditis; optic neuritis; chorioretinitis; anterior segment inflammation.

1.8 Respiratory Diseases:

Symptomatic sarcoidosis

1.9 Edematous State:

To induce a diuresis or a remission of proteinuria in the nephrotic syndrome without uremia of the idiopathic type or that due to lupus erythematosus.

2 DOSAGE AND ADMINISTRATION

2.1 Specific Recommended Dosage Regimen for Infantile Spasms in Infants and Children Under 2 Years of Age

In the treatment of infantile spasms, H.P. Acthar Gel must be administered intramuscularly. The recommended regimen is a daily dose of 150 U/m² (divided into twice daily intramuscular injections of 75 U/m²) administered over a 2-week period. Dosing with H.P. Acthar Gel should then be gradually tapered over a 2-week period to avoid adrenal insufficiency. The following is one suggested tapering schedule: 30 U/m² in the morning for 3 days; 15 U/m² in the morning for 3 days; 10 U/m² in the morning for 3 days; and 10 U/m² every other morning for 6-days.

H.P. Acthar Gel is typically dosed based on body surface area (BSA). For calculation of body surface area, use the following formula

$$BSA(m^2) = \sqrt{\frac{weight \text{ (kg)} \times height \text{ (cm)}}{3600}}$$

2.2 Recommended Dosage Regimen for the Treatment of Acute Exacerbations in Adults with Multiple Sclerosis.

The recommended dose is daily intramuscular or subcutaneous doses of 80-120 units for 2-3 weeks for acute exacerbations.

Dosage should be individualized according to the medical condition of each patient. Frequency and dose of the drug should be determined by considering the severity of the disease and the initial response of the patient.

Although drug dependence does not occur, sudden withdrawal of H.P. Acthar Gel after prolonged use may lead to adrenal insufficiency or recurrent symptoms which make it difficult to stop the treatment. It may be necessary to taper the dose and increase the injection interval to gradually discontinue the medication.

2.3 Recommended Dosage Regimen for Other Indications for Adults and Children Over 2 Years of Age

Dosage should be individualized according to the disease under treatment and the general medical condition of each patient. Frequency and dose of the drug should be determined by considering severity of the disease and the initial response of the patient.

The usual dose of H.P. Acthar Gel is 40-80 units given intramuscularly or subcutaneously every 24-72 hours.

Although drug dependence does not occur, sudden withdrawal of H.P. Acthar Gel after prolonged use may lead to adrenal insufficiency or recurrent symptoms which make it difficult to stop the treatment. It may be necessary to taper the dose and increase the injection interval to gradually discontinue the medication.

2.4 Preparation

H.P. Acthar Gel should be warmed to room temperature before using.

Caution should be taken not to over-pressurize the vial prior to withdrawing the product.

3 DOSAGE FORMS AND STRENGTHS

5 mL multi-dose vial containing 80 USP Units per mL.

4 CONTRAINDICATIONS

H.P. Acthar Gel is contraindicated for intravenous administration.

H.P. Acthar Gel is contraindicated where congenital infections are suspected in infants.

Administration of live or live attenuated vaccines is contraindicated in patients receiving immunosuppressive doses of H.P. Acthar Gel.

H.P. Acthar Gel is contraindicated in patients with scleroderma, osteoporosis, systemic fungal infections, ocular herpes simplex, recent surgery, history of or the presence of a peptic ulcer, congestive heart failure, uncontrolled hypertension, primary adrenocortical insufficiency, adrenocortical hyperfunction or sensitivity to proteins of porcine origin.

5 WARNINGS AND PRECAUTIONS

The adverse effects of H.P. Acthar Gel are related primarily to its steroidogenic effects. Not all of the adverse events described below have been seen after treatment with H.P. Acthar Gel, but might be expected to occur. [*see Adverse Reactions (6.3)*].

5.1 Infections

H.P. Acthar Gel may increase the risks related to infections with any pathogen, including viral, bacterial fungal, protozoan or helminthic infections. Patients with latent tuberculosis or tuberculin reactivity should be observed closely, and if therapy is prolonged, chemoprophylaxis should be instituted.

5.2 Cushing's Syndrome and Adrenal Insufficiency Upon Withdrawal

Treatment with H.P. Acthar Gel can cause hypothalamic-pituitary-axis (HPA) suppression and Cushing's syndrome. These conditions should be monitored especially with chronic use.

Suppression of the HPA may occur following prolonged therapy with the potential for adrenal insufficiency after withdrawal of the medication. Patients should be monitored for signs of insufficiency such as weakness, hyperpigmentation, weight loss, hypotension and abdominal pain.

The symptoms of adrenal insufficiency in infants treated for infantile spasms can be difficult to identify. The symptoms are non-specific and may include anorexia, fatigue, lethargy, weakness, excessive weight loss, hypotension and abdominal pain. It is critical that parents and caregivers be made aware of the possibility of adrenal insufficiency when discontinuing H.P. Acthar Gel and should be instructed to observe for, and be able to recognize, these symptoms [*see Information for Patients (17)*]

The recovery of the adrenal gland may take from days to months so patients should be protected from the stress (e.g. trauma or surgery) by the use of corticosteroids during the period of stress.

The adrenal insufficiency may be minimized in adults and infants by tapering of the dose when discontinuing treatment.

Signs or symptoms of Cushing's syndrome may occur during therapy but generally resolve after therapy is stopped. Patients should be monitored for these signs and symptoms such as deposition of adipose tissue in characteristic sites (e.g., moon face, truncal obesity), cutaneous striae, easy bruisability, decreased bone mineralization, weight gain, muscle weakness, hyperglycemia, and hypertension.

5.3 Elevated Blood Pressure, Salt and Water Retention and Hypokalemia

H.P. Acthar Gel can cause elevation of blood pressure, salt and water retention, and increased excretion of potassium and calcium. Dietary salt restriction and potassium supplementation may

be necessary. Caution should be used in the treatment of patients with hypertension, congestive heart failure, or renal insufficiency.

5.4 Vaccination

Administration of live or live attenuated vaccines is contraindicated in patients receiving immunosuppressive doses of H.P. Acthar Gel. Killed or inactivated vaccines may be administered; however, the response to such vaccines can not be predicted. Other immunization procedures should be undertaken with caution in patients who are receiving H.P. Acthar Gel, especially when high doses are administered, because of the possible hazards of neurological complications and lack of antibody response.

5.5 Masking Symptoms of Other Diseases

H.P. Acthar Gel often acts by masking symptoms of other diseases/disorders without altering the course of the other disease/disorder. Patients should be monitored carefully during and for a period following discontinuation of therapy for signs of infection, abnormal cardiac function, hypertension, hyperglycemia, change in body weight and fecal blood loss.

5.6 Gastrointestinal Perforation and Bleeding

H.P. Acthar Gel can cause GI bleeding and gastric ulcer. There is also an increased risk for perforation in patients with certain gastrointestinal disorders. Signs of gastrointestinal perforation, such as peritoneal irritation, may be masked by the therapy. Use caution where there is the possibility of impending perforation, abscess or other pyogenic infections, diverticulitis, fresh intestinal anastomoses, and active or latent peptic ulcer.

5.7 Behavioral and Mood Disturbances

Use of H.P. Acthar Gel may be associated with central nervous system effects ranging from euphoria, insomnia, irritability (especially in infants), mood swings, personality changes, and severe depression, to frank psychotic manifestations. Also, existing emotional instability or psychotic tendencies may be aggravated.

5.8 Comorbid Diseases

Patients with a comorbid disease may have that disease worsened. Caution should be used when prescribing H.P. Acthar Gel in patients with diabetes and myasthenia gravis.

5.9 Ophthalmic Effects

Prolonged use of H.P. Acthar Gel may produce posterior subcapsular cataracts, glaucoma with possible damage to the optic nerves and may enhance the establishment of secondary ocular infections due to fungi and viruses.

5.10 Immunogenicity Potential

H.P. Acthar Gel is immunogenic. Limited available data suggest that a patient may develop antibodies to H.P. Acthar Gel after chronic administration and loss of endogenous ACTH and H.P. Acthar Gel activity. Prolonged administration of H.P. Acthar Gel may increase the risk of hypersensitivity reactions. Sensitivity to porcine protein should be considered before starting therapy and during the course of treatment should symptoms arise.

5.11 Use in Patients with Hypothyroidism or Liver Cirrhosis

There is an enhanced effect in patients with hypothyroidism and in those with cirrhosis of the liver.

5.12 Negative Effects on Growth and Physical Development

Long-term use of H.P. Acthar Gel may have negative effects on growth and physical development in children. Changes in appetite are seen with H.P. Acthar Gel therapy, with the effects becoming more frequent as the dose or treatment period increases. These effects are reversible once H.P. Acthar Gel therapy is stopped. Growth and physical development of pediatric patients on prolonged therapy should be carefully monitored.

5.13 Decrease in Bone Density

Decrease in bone formation and an increase in bone resorption both through an effect on calcium regulation (i.e. decreasing absorption and increasing excretion) and inhibition of osteoblast function may occur. These, together with a decrease in the protein matrix of the bone (secondary to an increase in protein catabolism) and reduced sex hormone production, may lead to inhibition of bone growth in children and adolescents and to the development of osteoporosis at any age. Special consideration should be given to patients at increased risk of osteoporosis (i.e., postmenopausal women) before initiating therapy, and bone density should be monitored in patients on long term therapy.

5.14 Use in Pregnancy

H.P. Acthar Gel has been shown to have an embryocidal effect. Apprise women of potential harm to the fetus. [*see Use in Specific Populations (8.1)*]

6 ADVERSE REACTIONS

Please refer to *Adverse Reactions in Infants and Children Under 2 Years of Age (Section 6.1.1)* for consideration when treating patients with Infantile Spasms. The adverse reactions presented in Section 6.2 are primarily provided for consideration in use in adults and in children over 2 years of age, but these adverse reactions should also be considered when treating infants and children under 2 years of age.

H.P. Acthar Gel causes the release of endogenous cortisol from the adrenal gland. Therefore all the adverse effects known to occur with elevated cortisol may occur with H.P. Acthar Gel administration as well. Common adverse reactions include fluid retention, alteration in glucose tolerance, elevation in blood pressure, behavioral and mood changes, increased appetite and weight gain.

6.1 Clinical Studies Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug, and may not reflect the rates observed in practice.

6.1.1 Adverse Reactions in Infants and Children Under 2 Years of Age

While the types of adverse reactions seen in infants and children under age 2 treated for infantile spasms are similar to those seen in older patients, their frequency and severity may be different due to the very young age of the infant, the underlying disorder, the duration of therapy and the dosage regimen. Below is a summary of adverse reactions specifically tabulated from source data derived from retrospective chart reviews and clinical trials in children under 2 years of age treated for infantile spasms. The number of patients in controlled trials at the recommended dose was too few to provide meaningful incidence rates or to permit a meaningful comparison to the control groups.

TABLE: Incidence (%) of Treatment Emergent Adverse Events Occurring in $\geq 2\%$ of H.P. Acthar Gel (repository corticotropin injection) Infants and Children under 2 years of Age

System Organ Class	Recommended 75 U/m ² bid n=122, (%)	150 U/m ² qd n=37 (%)
Cardiac disorders		
Cardiac Hypertrophy	3	0
Endocrine disorders		
Cushingoid	3	22
Gastrointestinal disorders		
Constipation	0	5
Diarrhea	3	14
Vomiting	3	5
General disorders and administration site conditions		
Irritability	7	19
Pyrexia	5	8
Infections and infestations		
Infection ¹	20	46

System Organ Class	Recommended 75 U/m² bid n=122, (%)	150 U/m² qd n=37 (%)
Investigations		
Weight gain	1	3
Metabolism and nutrition disorders		
Increased appetite	0	5
Decreased appetite	3	3
Nervous system disorders		
Convulsion ²	12	3
Respiratory, thoracic and mediastinal disorders		
Nasal Congestion	1	5
Skin and subcutaneous tissue disorders		
Acne	0	14
Rash	0	8
Vascular disorders		
Hypertension	11	19

¹ Specific infections that occurred at $\geq 2\%$ were candidiasis, otitis media, pneumonia and upper respiratory tract infections. ² In the treatment of Infantile Spasms, other types of seizures/convulsions may occur because some patients with infantile spasms progress to other forms of seizures (for example, Lennox-Gastaut Syndrome). Additionally the spasms sometimes mask other seizures and once the spasms resolve after treatment, the other seizures may become visible.

These adverse reactions may also be seen in adults and children over 2 years of age when treated for other purposes and with different doses and regimens.

6.2 Postmarketing Experience

The following adverse reactions associated with the use of H.P. Acthar Gel have been identified from postmarketing experience with H.P. Acthar Gel. Only adverse events that are not listed above as adverse events reported from retrospective chart reviews and non-sponsor conducted clinical trials and those not discussed elsewhere in labeling, are listed in this section. Because the adverse reactions are reported voluntarily from a population of uncertain size, it is not always possible to estimate their frequency or establish a causal relationship to use with H.P. Acthar Gel. Events are categorized by system organ class. Unless otherwise noted these adverse events have been reported in infants, children and adults.

6.2.1 Allergic Reactions

Allergic responses have presented as dizziness, nausea and shock (adults only).

6.2.2 Cardiovascular

Necrotizing angitis (adults only) and congestive heart failure.

6.2.3 Dermatologic

Skin thinning (adults only), facial erythema and increased sweating (adults only).

6.2.4 Endocrine

Decreased carbohydrate tolerance (infants only) and hirsutism.

6.2.5 Gastrointestinal

Pancreatitis (adults only), abdominal distention and ulcerative esophagitis.

6.2.6 Metabolic

Hypokalemic alkalosis (infants only).

6.2.7 Musculoskeletal

Muscle weakness and vertebral compression fractures (infants only).

6.2.8 Neurological

Headache (adults only), vertigo (adults only), subdural hematoma, intracranial hemorrhage (adults only), and reversible brain shrinkage (usually secondary to hypertension) (infants only).

6.3 Possible Additional Steroidogenic Effects

Based on steroidogenic effects of H.P. Acthar Gel certain adverse events may be expected due to the pharmacological effects of corticosteroids. The adverse events that may occur but have not been reported for H.P. Acthar Gel are:

6.3.1 Dermatologic

Impaired wound healing, abscess, petechiae and ecchymoses, and suppression of skin test reactions.

6.3.2 Endocrine

Menstrual irregularities.

6.3.3 Metabolic

Negative nitrogen balance due to protein catabolism.

6.3.4 Musculoskeletal

Loss of muscle mass and aseptic necrosis of femoral and humeral heads.

6.3.5 Neurological

Increased intracranial pressure with papilledema, (pseudo-tumor cerebri) usually after treatment, and subdural effusion.

6.3.6 Ophthalmic

Exophthalmos.

7 DRUG INTERACTIONS

Formal drug-drug interaction studies have not been performed.

H.P. Acthar Gel may accentuate the electrolyte loss associated with diuretic therapy.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Class C: H.P. Acthar Gel has been shown to have an embryocidal effect. There are no adequate and well-controlled studies in pregnant women. H.P. Acthar Gel should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

8.3 Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from H.P. Acthar Gel, when treating a nursing mother, a decision should be made whether to discontinue nursing or to discontinue the drug, considering the risk and benefit to the mother.

8.4 Pediatric Use

H.P. Acthar Gel is indicated as monotherapy for the treatment of infantile spasms in infants and children less than 2 years of age. Both serious and other adverse reactions in this population are discussed in Warnings and Adverse Reactions in Infants and Children Under 2 Years of Age [*see Sections 5 and 6.1.1*].

The efficacy of H.P. Acthar Gel for the treatment of infantile spasms in infants and children less than 2 years of age was evaluated in a randomized, single blinded (video EEG interpreter

blinded) clinical trial and an additional active control supportive trial [see *Clinical Studies (14)*]. A responding patient was defined as having both complete cessation of spasms and elimination of hypsarrhythmia.

Safety in the pediatric population for infantile spasms was evaluated by retrospective chart reviews and data from non-sponsor conducted clinical trials [see *Adverse Reactions (6.1.1)*]. While the types of adverse reactions seen in infants and children under 2 years of age treated for infantile spasms are similar to those seen in older patients, their frequency and severity may be different due to the very young age of the infant, the underlying disorder, the duration of therapy and the dosage regimen. Effects on growth are of particular concern [see *Warnings and Precautions (5.12)*]. Serious adverse reactions observed in adults may also occur in children [see *Warnings and Precautions (5)*].

10 OVERDOSAGE

While chronic exposure to H.P. Acthar Gel at high doses can be associated with a variety of potential serious adverse effects, it is not expected that a single high dose, or even several large doses, has the potential for serious adverse effects compared to a standard dose. There have been no reports of death or acute overdose symptoms from H.P. Acthar Gel in clinical studies or in the published literature.

The intramuscular route of administration makes it unlikely that an inadvertent acute overdose will occur. The typical daily dose of H.P. Acthar Gel to treat an infant that has a BSA of 0.4 m² would be 60 U/day. Using the 1-cc syringe supplied with H.P. Acthar Gel, the maximum amount that can be injected is 80 U/injection, which is a well-tolerated single dose.

11 DESCRIPTION

H.P. Acthar Gel is a highly purified sterile preparation of the adrenocorticotrophic hormone in 16% gelatin to provide a prolonged release after intramuscular or subcutaneous injection. Also contains 0.5% phenol, not more than 0.1% cysteine (added), sodium hydroxide and/or acetic acid to adjust pH and water for injection.

ACTH is a 39 amino acid peptide with the following chemical formula:

H-	Ser-	Tyr-	Ser-	Met-	Glu-	His-	Phe-	Arg-	Trp-	Gly-
	1	2	3	4	5	6	7	8	9	10
	Lys-	Pro-	Val-	Gly-	Lys-	Lys-	Arg-	Arg-	Pro-	Val-
	11	12	13	14	15	16	17	18	19	20
	Lys-	Val-	Try-	Pro-	Asp-	Gly-	Ala-	Glu-	Asp-	Gln-
	21	22	23	24	25	26	27	28	29	30
	Leu-	Ala-	Glu-	Ala-	Phe-	Pro-	Leu-	Glu-	Phe-	OH
	31	32	33	34	35	36	37	38	39	

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism of action of H.P. Acthar Gel in the treatment of infantile spasms is unknown.

H.P. Acthar Gel and endogenous ACTH stimulate the adrenal cortex to secrete cortisol, corticosterone, aldosterone, and a number of weakly androgenic substances. Prolonged administration of large doses of H.P. Acthar Gel induces hyperplasia and hypertrophy of the adrenal cortex and continuous high output of cortisol, corticosterone and weak androgens. The release of endogenous ACTH is under the influence of the nervous system via the regulatory hormone released from the hypothalamus and by a negative corticosteroid feedback mechanism. Elevated plasma cortisol suppresses ACTH release.

H.P. Acthar Gel is also reported to bind to melanocortin receptors.

The trophic effects of endogenous ACTH and H.P. Acthar Gel on the adrenal cortex are not well understood beyond the fact that they appear to be mediated by cyclic AMP.

ACTH rapidly disappears from the circulation following its intravenous administration; in people, the plasma half-life is about 15 minutes. The pharmacokinetics of H.P. Acthar Gel have not been adequately characterized.

The maximal effects of a trophic hormone on a target organ are achieved when optimal amounts of hormone are acting continuously. Thus, a fixed dose of H.P. Acthar Gel will demonstrate a linear increase in adrenocortical secretion with increasing duration for the infusion.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Adequate and well-controlled studies have not been done in animals. Human use has not been associated with an increase in malignant disease. [*see Warnings and Precautions (5.14) and Use in Specific Populations (8.1)*].

14 CLINICAL STUDIES

The effectiveness of H.P. Acthar Gel as a treatment for infantile spasms was demonstrated in a single blinded (video EEG interpreter blinded) clinical trial in which patients were randomized to receive either a 2 week course of treatment with H.P. Acthar Gel (75 U/m² intramuscular twice daily) or prednisone (1 mg/kg by mouth twice daily). The primary outcome was a comparison of the number of patients in each group who were treatment responders, defined as a patient having complete suppression of both clinical spasms and hypsarrhythmia on a full sleep cycle video EEG performed 2 weeks following treatment initiation, rated by an investigator blinded to treatment. Thirteen of 15 patients (86.7%) responded to H.P. Acthar Gel as compared to 4 of 14

patients (28.6%) given prednisone ($p < 0.002$). The 2-week treatment was followed by a 2-week period of taper. Nonresponders to the prednisone treatment were eligible to receive H.P. Acthar Gel treatment. Seven of 8 patients (87.5%) responded to H.P. Acthar Gel after not responding to prednisone. Similarly, the 2 nonresponder patients from the H.P. Acthar Gel treatment were eligible to receive treatment with prednisone. One of the 2 patients (50%) responded to the prednisone treatment after not responding to H.P. Acthar Gel.

A supportive single-blind, randomized clinical trial comparing high-dose, long-duration treatment (150 U/m² once daily for 3 weeks, n=30) of H.P. Acthar Gel with low-dose, short-duration treatment (20 U once daily for 2 weeks, n=29) for the treatment of infantile spasms was also evaluated in infants and children less than 2 years of age. Nonresponders (defined as in the previously described study) in the low-dose group received a dose escalation at 2 weeks to 30 U once daily. Nominal statistical superiority of the high dose treatment, as compared to the low dose treatment, was observed for cessation of spasms but not for the resolution of hypsarrhythmia.

16 HOW SUPPLIED / STORAGE AND HANDLING

H.P. Acthar Gel (repository corticotropin injection) is supplied as 5 mL multi-dose vial (63004-7731-1) containing 80 USP Units per mL. H.P. Acthar Gel (repository corticotropin injection) should be warmed to room temperature before using. Do not over pressurize the vial prior to withdrawing the product.

Store H.P. Acthar Gel (repository corticotropin injection) under refrigeration between 2°-8°C (36°-46°F). Product is stable for the period indicated on the label when stored under the conditions described.

17 PATIENT COUNSELING INFORMATION

Caretakers of patients with infantile spasms should be informed of the availability of a Medication Guide, and they should be instructed to read the Medication Guide prior to administering H.P. Acthar Gel. Patients should be instructed to take H.P. Acthar Gel only as prescribed. They should not stop treatment suddenly unless instructed by their physician to do so.

Patients, their caregivers and families should be advised as to the importance of the need for careful monitoring while on and during titration from H.P. Acthar Gel treatment and the importance of not missing and scheduled doctor's appointments.

Patients, their caregivers and families should be advised that if the patient develops an infection or fever they should contact their physician. They should be educated that a fever may not necessarily be present during infection. The patient should also try to limit contact with other people with infections to minimize the risk of infection while taking H.P. Acthar Gel. [*see Warnings and Precautions (5.1) and Adverse Reactions (6.1.1)*]

Patients, their caregivers and families should be advised that if the patient experiences an increase in blood pressure they should contact their physician. [*see Warnings and Precautions (5.3) and Adverse Reactions (6.1.1)*]

Patients, their caregivers and families should be advised that if the patient or the caregiver notices blood or a change in color of the patient's stool they should contact their physician. [*see Warnings and Precautions (5.6)*].

Caregivers and families of infants and children treated with H.P. Acthar Gel should be informed that the patient may show signs of irritability and sleep disturbances. These effects are reversible once H.P. Acthar Gel therapy is stopped. [*see Warnings and Precautions (5.7) and Adverse Reactions (6.1.1)*].

Patients, their caregivers and families should be advised that changes in appetite, most often leading to weight gain, are seen with H.P. Acthar Gel therapy, becoming more frequent as the dose or treatment period increases. These effects are reversible once H.P. Acthar Gel therapy is stopped. [*see Warnings and Precautions (5.12) and Adverse Reactions (6.1.1)*].

Patients, their caregivers and families should be advised that the patient may be monitored for signs of adrenal insufficiency such as weakness, fatigue, lethargy, anorexia, weight loss, hypotension, abdominal pain or hyperpigmentation (adults only) after treatment has stopped. Since the recovery of the adrenal gland varies from days to months, patients may need to be protected from the stress of trauma or surgery by the use of corticosteroids during the period of stress. [*see Warnings and Precautions (5.2)*].

Patients should be advised not to be vaccinated with live or live attenuated vaccines during treatment with H.P. Acthar Gel. Additionally, other immunization procedures in patients or in family members who will be in contact with the patient should be undertaken with caution while the patient is taking H.P. Acthar Gel. [*see Warnings and Precautions (5.4)*].

Patients, their caregivers and families should be advised that prolonged use of H.P. Acthar Gel in children may result in Cushing's syndrome and associated adverse reactions, may inhibit skeletal growth, and may cause osteoporosis and decreased bone density. If prolonged use is necessary, H.P. Acthar Gel should be given intermittently along with careful observation. [*see Warnings and Precautions (5.2), (5.12), and (5.13) and Adverse Reactions (6.1.1)*].

Patients, their caregivers and families should be informed that H.P. Gel may mask symptoms of other diseases/disorders without altering the course of the other disease/disorder. The patient will need to be monitored carefully during and for a period following discontinuation of therapy for signs of infection, abnormal cardiac function, hypertension, hyperglycemia, change in body weight, and fecal blood loss. [*see Warnings and Precautions (5.5)*].

In the treatment of Infantile Spasms, other types of seizures may occur because some patients with infantile spasms progress to other forms of seizures (for example, Lennox-Gastaut Syndrome). Additionally the spasms sometimes mask other seizures and once the spasms resolve after treatment with H.P. Acthar Gel, the other seizures may become visible. Parents and caregivers should inform their physician of any new onset of seizures so that appropriate management can then be instituted. [*see Adverse Reactions (6.1.1)*].

H.P. Acthar[®] Gel
(repository corticotropin injection)

Manufactured for Questcor Pharmaceuticals, Inc.



QUESTCOR[®]

Questcor Pharmaceuticals, Inc.

3260 Whipple Road

Union City, CA 94587 USA

phone (800) 411-3065

(510) 400-0700

fax (510) 400-0799

PL065/Rev. 02

No. 1350

Issued: 10/2010

PM-210-03 10/10